

Effect of Interaction between Hepatitis C Virus NS5A and NS5B on Hepatitis C Virus RNA Replication with the Hepatitis C Virus Replicon

Tetsuro Shimakami,^{1,2} Makoto Hijikata,³ Hong Luo,^{1,4} Yuan Yuan Ma,¹
Shuichi Kaneko,² Kunitada Shimotohno,³ and Seishi Murakami^{1*}

Department of Molecular Oncology, Cancer Research Institute, Kanazawa University,¹ and Department of Gastroenterology, Kanazawa University Graduate School of Medicine,² Takara-Machi, Kanazawa, Ishikawa 920-0934, and Department of Viral Oncology, Institute for Virus Research, Kyoto University, Sakyo-Ku, Kyoto 606-8507,³ Japan, and The Institute of Infectious Diseases, West China Hospital, Sichuan University, Chengdu, Sichuan 610041, China⁴

Received 3 September 2003/Accepted 10 November 2003

Hepatitis C virus (HCV) NS5A has been reported to be important for the establishment of replication by adaptive mutations or localization, although its role in viral replication remains unclear. It was previously reported that NS5A interacts with NS5B via two regions of NS5A in the isolate JK-1 and modulates the activity of NS5B RdRp (Y. Shiota et al., *J. Biol. Chem.*, 277:11149–11155, 2002), but the biological significance of this interaction has not been determined. In this study, we addressed the effect of this interaction on HCV RNA replication with an HCV replicon system derived from the isolate M1LE (H. Kishine et al., *Biochem. Biophys. Res. Commun.*, 293:993–999, 2002). We constructed three internal deletion mutants, M1LE/5Adel-1 and M1LE/5Adel-2, each encoding NS5A which cannot bind NS5B, and M1LE/5Adel-3, encoding NS5A that can bind NS5B. After transfection into Huh-7 cells, M1LE/5Adel-3 was replication competent, but both M1LE/5Adel-1 and M1LE/5Adel-2 were not. Next we prepared 20 alanine-substituted clustered mutants within both NS5B-binding regions and examined the effect of these mutants on HCV RNA replication. Only 5 of the 20 mutants were replication competent. Subsequently, we introduced a point mutation, S225P, a deletion of S229, or S232I into NS5A and prepared cured Huh-7 cells that were cured of RNA replication by alpha interferon. Finally, with these point mutations and cured cells, we established a highly improved replicon system. In this system, only the same five mutants were replication competent. These results strongly suggest that the interaction between NS5A and NS5B is critical for HCV RNA replication in the HCV replicon system.

The hepatitis C virus (HCV) is a major cause of chronic hepatitis around the world (1, 9). Chronic infection with HCV results in liver cirrhosis and often hepatocellular carcinoma (50, 53). HCV is an enveloped positive-strand RNA virus belonging to the genus *Hepacivirus* in the family *Flaviviridae* (44). The HCV RNA genome is ~9.6 kb in length and consists of a 5' nontranslated region (5' NTR), a large open reading frame, and a 3' NTR. The 5' NTR contains an internal ribosome entry site (IRES) mediating translation of a single polypeptide of ~3,000 amino acid (aa) residues (57, 59). The polypeptide is cleaved by host and viral protease into at least 10 different products (2, 21, 22, 25, 26). The structural proteins core, E1, and E2 are located in the amino terminus of the polypeptide, followed by p7, a hydrophobic peptide with unknown function, and the nonstructural (NS) proteins NS2, NS3, NS4A, NS4B, NS5A, and NS5B (41). The 3' NTR consists of a short variable sequence, a poly(U)-poly(UC) tract, and a highly conserved X region, and it is critical for HCV RNA replication and HCV infection (17, 34, 65, 66).

HCV is unique among positive-strand RNA viruses in causing persistent infections, and a high mutation rate in E2 allows

it to escape host immune surveillance. These phenomena are tightly associated with chronic inflammation of the liver (27, 32, 61, 62). Therefore, HCV RNA replication has been a target for treatment of HCV. NS5B is an RNA-dependent RNA polymerase (RdRp), the central catalytic enzyme in HCV RNA replication. Several recombinant forms of NS5B expressed and purified from insect cells and *Escherichia coli* are available and catalytically active, and studies with purified NS5B proteins provide insight into the biochemical and catalytic properties of NS5B (3, 16, 39, 64). However, the result that NS5B can initiate de novo RNA replication in vitro with both a non-HCV RNA template and an HCV RNA template may reflect the catalytic property of NS5B but not the tight regulation of HCV RNA replication initiation. Studies of HCV RNA replication in vitro have to overcome several difficulties, since replication requires all or most NS proteins and occurs at the membrane where all of the HCV NS proteins are recruited.

A second system used to study HCV RNA replication is the study of HCV RNA replicons in vivo, which utilizes autonomously replicating HCV-derived RNAs. These replicon RNAs have the authentic HCV 5' and 3' NTRs. The HCV IRES drives the translation of a selectable marker such as neomycin resistance, and an internal encephalomyocarditis virus IRES directs translation of NS3 to NS5B (38). In vitro-transcribed replicon RNAs are transfected into the human hepatoma cell line Huh-7 by electroporation and placed under selection. The emergence of neomycin-resistant cell colonies is indicative of

* Corresponding author. Mailing address: Department of Molecular Oncology, Cancer Research Institute, Kanazawa University, 13-1 Takara-Machi, Kanazawa, 920-0934 Ishikawa, Japan. Phone: 81-76-265-2731. Fax: 81-76-234-4501. E-mail: semuraka@kenroku.kanazawa-u.ac.jp.

RNA replication. Recently, adaptive mutations that dramatically enhance the ability of HCV RNA to replicate have been identified in NS3, NS4B, NS5A, and NS5B (4, 24, 35, 37, 40). Initially, only the replicon derived from a genotype 1b isolate, HCV-Con1, was replication competent in Huh-7 cells, and adaptive mutations were required for efficient replication (4, 35, 37, 40). Then, several replicon systems derived from a genotype 1b isolate, HCV-N, and a genotype 1a isolate, HCV-H77, were reported to replicate in Huh-7 cells (6, 23, 24, 29). Recently, a replicon system was established which uses an HCV sequence derived from the human T-cell line MT-2C infected with HCV (a genotype 1b isolate, M1LE) in vitro and isolated 50-1 cells replicating subgenomic RNAs with some amino acid mutations (31, 33, 43, 55).

It was previously reported that NS5A and NS5B interact in vitro and in vivo through two independent regions of NS5A and that NS5A modulates the activity of NS5B RdRp through this interaction in vitro in the isolate JK-1 (54). This ability of NS5A to modulate the RdRp activity in vitro may be consistent with the high frequency of adaptive mutations in NS5A which result in a much more efficient RNA replication in Huh-7 cells; however, the biological meaning of the interaction between NS5A and NS5B remains to be addressed in vivo. Here we report a modified HCV RNA replicon system derived from the isolate M1LE by introducing point mutations (S225P, a deletion of S229, and S232I), and curing 50-1 cells of HCV subgenomic RNA replication by interferon (IFN) treatment. By introducing internal deletion and substitution mutations into NS5A, we demonstrate that the regions essential for the interaction between NS5A and NS5B are also critical for HCV RNA replication in using the HCV replicon system.

MATERIALS AND METHODS

Construction of plasmids. pNNRZ2RU (33) harbors a subgenomic replicon derived from MT-2C cells infected with HCV (a genotype 1b isolate, M1LE; GenBank accession no. AB080299), and this plasmid contains cDNA of wild-type M1LE. For convenience, pNNRZ2RU was digested with *Mlu*I and *Bgl*II, and the obtained fragment was inserted into the *Mlu*I and *Bgl*II sites of the vector pGL3Basic (Promega) to create pGL3-*Mlu*I-*Bgl*II. pGL3-*Mlu*I-*Bgl*II was used as an intermediate vector. All mutations were introduced into pGL3-*Mlu*I-*Bgl*II, and then the fragments of pGL3-*Mlu*I-*Bgl*II digested by *Mlu*I and *Bgl*II containing each mutation were reintroduced into the *Mlu*I and *Bgl*II sites of pNNRZ2RU to create each mutant.

M1LE/5Adel-1 was generated by PCR with the primers 5Adel-1 For (containing a *Mlu*I site) and 5A Rev-3 and then inserted into the *Mlu*I and *Nhe*I sites of pGL3-*Mlu*I-*Bgl*II to create pGL3-*Mlu*I-*Bgl*II-5Adel-1. M1LE/5Adel-2 was generated by PCR with overlap extension with the primers 5Adel-2 For, 5A Rev-2, 5Adel-2 Rev, and 5A For-3 and then inserted into the *Nhe*I and *Sac*I sites of pGL3-*Mlu*I-*Bgl*II to create pGL3-*Mlu*I-*Bgl*II-5Adel-2. M1LE/5Adel-3 was generated by PCR with the primers 5A del-3 For (containing a *Nhe*I site) and 5A Rev-2 and then inserted into the *Nhe*I and *Sac*I sites of pGL3-*Mlu*I-*Bgl*II to create pGL3-*Mlu*I-*Bgl*II-5Adel-3.

An alanine scanning method was used to construct NS5A alanine-substituted mutants to minimize the effects of substituted amino acid residues (7). The positions of alanine-substituted clustered mutations (cm) of NS5A are shown in Fig. 2B. To generate M1LE/cm 94, 100, 105, 110, 113, 120, 127, 134, 141, 148, and 155, each mutation was introduced into the *Mlu*I and *Nhe*I sites of pGL3-*Mlu*I-*Bgl*II by site-directed mutagenesis with primers carrying the necessary nucleotide changes to create pGL3-*Mlu*I-*Bgl*II-cm94, -100, -105, -110, -113, -120, -127, -134, -141, -148, and -155, respectively. To generate M1LE/cm 252, 277, 283, 290, 297, 304, 311, 316, 321, and 328, each mutation was introduced into the *Nhe*I and *Sac*I sites of pGL3-*Mlu*I-*Bgl*II by site-directed mutagenesis with primers carrying the necessary nucleotide changes to create pGL3-*Mlu*I-*Bgl*II-cm252, -277, -283, -290, -297, -304, -311, -316, -321, and -328, respectively.

To generate M1LE/S225P and a deletion of S229 (delS229), the point muta-

TABLE 1. Sequences of primers used in the present study

Primer	Sequence (5'-3')
5Adel-1 For.....	ATATATCAACGCGTACCCGGCGTGTAACCTCTCCTACGG
5Adel-2 For.....	GTGGAGTCAGAACGTTCTCCGGTGGTACACGGGTGCCCA
5Adel-2 Rev	TACCACCGGAGGAACGTTCTCTGACTCCACGCGGGTGTGTT
5Adel-3 For.....	ATATATATGCTAGCCAGTTGAAGGTAGTATTCTGGACTTTTC
5A For-3	ATCCTTCCCACATTACAGCA
5A Rev-2	CTCAACGTCGGATCCCTTGT
5A Rev-3	GGTCAGCGTCCGGGGAGTCATG
NS5A For	ATATCAATTGCATGTCCGGCTCGTGGCTAAGGATATT
NS5A Rev	ATATAGATCTGCAGCAGACGACGTCCTCACTAGCCTC
NS5B For	TATCGAGTCGATGTCAATGTCCTACTCATGGACAGGT
NS5B Rev.....	ATATGGGATCCCCGGTTGGGGAGCAGGTAGATGCTAC

tions S225P and delS229 were introduced into the *Mlu*I and *Nhe*I sites of pGL3-*Mlu*I-*Bgl*II by site-directed mutagenesis with primers carrying the necessary nucleotide changes to create pGL3-*Mlu*I-*Bgl*II-S225P and -delS229. The point mutation S232I was introduced into the *Mlu*I and *Sac*I sites of pGL3-*Mlu*I-*Bgl*II by site-directed mutagenesis with primers carrying the necessary nucleotide changes to create pGL3-*Mlu*I-*Bgl*II-S232I. To generate the double mutants containing both the point mutation S232I plus an internal deletion mutation or cm, the *Eco*RI fragments of pGL3-*Mlu*I-*Bgl*II-cm94, -100, -105, -110, -113, -120, -127, -134, and -141 were inserted into the *Eco*RI sites of pGL3-*Mlu*I-*Bgl*II-S232I. Because there were no optimal enzyme sites in the others, the point mutation S232I was introduced into the *Mlu*I and *Sac*I sites of pGL3-*Mlu*I-*Bgl*II-5Adel-1, -del-2, -del-3, and -cm148, -155, -252, -277, -283, -290, -297, -304, -311, -316, -321, and -328 by site-directed mutagenesis with primers carrying the necessary nucleotide changes.

To create double mutants containing both the point mutation S225P plus an internal deletion mutation or cm (cm 252, 277, 283, 290, 297, 304, 311, 316, 321, and 328), the *Nhe*I and *Sac*I fragments of pGL3-*Mlu*I-*Bgl*II-5Adel-2, -5Adel-3, -cm252, -277, -283, -290, -297, -304, -311, -316, -321, and -328 were introduced into the *Nhe*I and *Sac*I sites of pGL3-*Mlu*I-*Bgl*II-S225P.

To generate M1LE/5B-VDD, a point mutation changing the GDD motif of NS5B to VDD was introduced at the *Nde*I and *Sma*I sites of pGL3-*Mlu*I-*Bgl*II by site-directed mutagenesis with primers carrying the necessary nucleotide changes to create pGL3-*Mlu*I-*Bgl*II-5B-VDD.

All of the mammalian expression vectors were derived from pSG5UTPL (36). The vector pNKFLAG (49) was used to express amino-terminally FLAG-tagged proteins. The vector pNKGST (49) was used to express glutathione *S*-transferase (GST)-fused proteins.

pNNRZ2RU was subcloned by PCR with the primers NS5A For, NS5A Rev, NS5B For, and NS5B Rev. NS5A For contains an artificial initiation codon downstream of the *Mun*I site, and NS5B For contains one artificial initiation codon downstream of the *Sac*I site. NS5A Rev contains a *Bgl*II site, and NS5B Rev contains a *Bam*HI site. Full-length NS5A was subcloned into the *Eco*RI and *Bam*HI sites of pNKFLAG to create pNKFLAG-5A/wild, and full-length NS5B was subcloned into the *Sac*I and *Bam*HI sites of pNKGST to create pNKGST-5B/wild.

To create NKFLAG-5A/del-1, and -cm94, -100, -105, -113, -120, -127, -134, -141, -148, and -155, the *Mlu*I and *Nhe*I fragments of pGL3-*Mlu*I-*Bgl*II mutants were introduced at the *Mlu*I and *Nhe*I sites of pNKFLAG-5A/wild. To create NKFLAG-5A/del-2, -del-3, -cm252, -277, -283, -290, -297, -304, -311, -316, -321, and -328, the *Nhe*I and *Sac*I fragments of pGL3-*Mlu*I-*Bgl*II mutants were inserted into the *Nhe*I and *Sac*I sites of pNKFLAG-5A/wild.

The sequences of all of the constructs were confirmed by the dideoxy sequence method. The main primers used for plasmid construction are shown in Table 1.

In vitro transcription and purification of RNA. Plasmids were linearized with *Xba*I and purified by passage through a column (PCR purification kit; Qiagen) prior to transcription. RNA was synthesized with T7 MEGAScript reagents (Ambion) by following the manufacturer's directions, and the reaction was stopped by digestion with RNase-free DNase. The synthesized RNA was passed

through a column (RNeasy mini kit; Qiagen) and dissolved in RNase-free water. The concentration was determined by measuring the optical density at 260 nm, and RNA integrity was checked by nondenaturing agarose gel electrophoresis.

RNA transfection and selection of G418-resistant cells. Subconfluent Huh-7 cells were trypsinized, washed once with phosphate-buffered saline (PBS) (–), and resuspended at 10^7 cells/ml in OPTI-MEM (Gibco-BRL, Invitrogen Life Technologies). Then, 10 to 1,000 ng of transcript was adjusted with total RNA from naive Huh-7 cells to a final amount of 10 μ g, which was mixed with 400 μ l of the cell suspension in a cuvette with a gap width of 0.4 cm (Bio-Rad). The mixture was immediately transfected into Huh-7 cells by electroporation with GenePulser II system (Bio-Rad) set to 270 V and 975 μ F. Following 10 min of incubation at room temperature, the cells were transferred into 10 ml of growth medium and then seeded into a 10- or 15-cm-diameter cell culture dish. For the selection of G418-resistant cells, the medium was replaced with fresh medium containing 0.5 to 1 mg of G418 (Geneticin; Gibco-BRL, Invitrogen Life Technologies)/ml after 24 to 48 h and the medium was changed twice a week. Four weeks after transfection, colonies were stained with Coomassie brilliant blue (0.6 g/liter in 50% methanol–10% acetic acid).

IFN treatment. To stop the replication of HCV subgenomic RNA, 50-1 cells were treated with 10,000 U of IFN- α 2b (kindly provided by Schering-Plough)/ml in the absence of G418. After 2 weeks of IFN treatment, the absence of HCV RNA was determined from the results of Northern hybridization, reverse transcription-PCR, and sensitivity to G418.

Cell culture. We used two kinds of Huh-7 cells, one derived from our own laboratory's original Huh-7 cell line, designated Huh-7-DMB, and the cured clone of 50-1 cells, designated Huh-7-KV-C. Both types of Huh-7 cells were grown in Dulbecco's modified Eagle's medium (Gibco-BRL, Invitrogen Life Technologies) supplemented with 10% fetal bovine serum, 2 mM L-glutamine, nonessential amino acids, 100 U of penicillin, and 100 μ g of streptomycin.

Preparation of cell extracts, coprecipitation with glutathione resin, and Western blot analysis. The transient transfection of COS1 cells was carried out by using the calcium-phosphate method. The cells were harvested, washed with PBS (–), and sonicated in PBS lysis buffer [PBS (–) containing 250 mM NaCl, 1.0% Triton X-100, 1 mM EDTA, and 1 mM dithiothreitol] with 10 mg (each) of aprotinin and leupeptin per ml. Total cell lysate was diluted 10-fold with PBS lysis buffer, mixed with 40 μ l of glutathione-Sepharose 4B beads (glutathione resin) (Amersham Biosciences), and then incubated for 3 h on a rotator in a cold room. After an extensive wash with PBS (–) containing 1.0% Triton X-100, the bound proteins were eluted, fractionated by sodium dodecyl sulfate (SDS)–10% polyacrylamide gel electrophoresis (PAGE), transferred onto nitrocellulose membranes, and subjected to Western blot analysis with anti-FLAG monoclonal antibody. The proteins were visualized by enhanced chemiluminescence according to the manufacturer's instructions (Amersham Biosciences). The nitrocellulose membranes used for Western blot analysis with anti-FLAG monoclonal antibody were reprobed with anti-GST monoclonal antibody (Zymed Laboratories) according to the manufacturer's instructions (Amersham Biosciences).

RESULTS

Interaction between NS5A and NS5B. It was previously reported that NS5A and NS5B associate through two discontinuous regions of NS5A (aa 105 to 162 and 277 to 334) and that NS5A weakly stimulates the activity of NS5B RdRp in vitro initially (at a molar ratio to NS5B of less than 0.1) and then inhibits the activity in a dose-dependent manner (54). To examine the effect of this interaction on HCV RNA replication, we used an HCV RNA replicon system derived from the isolate M1LE and 50-1 cells (33). We prepared a hybrid replicon of JK-1 and M1LE which harbors the JK-1 sequence from aa 92 (*Mlu*I site) of NS5A to the end of NS5B. However, the hybrid replicon did not produce any G418-resistant colony with the Huh-7 cell line (data not shown), so we constructed various mutated versions of NS5A of the RNA replicon derived from M1LE.

First, we confirmed whether the association between NS5A and NS5B through the two discontinuous regions of NS5A occurs with the sequence derived from M1LE. COS1 cells were transiently cotransfected with mammalian expression vectors,

pNKFLAG-5A/wild, /del-1, /del-2, and /del-3; pNKGST or pNKGST-5B/wild; and the cell lysates were subjected to a GST pull-down assay. pNKFLAG-5A/wild encodes the full-length and wild-type NS5A proteins of M1LE. pNKFLAG-5A/del-1 encodes the internally deleted NS5A protein missing aa 94 to 162, a deletion 11 aa longer than that reported for JK-1 for the convenience of mutagenesis (aa 105 to 162). pNKFLAG-5A/del-2 encodes the internally deleted NS5A protein missing aa 277 to 334, the same region reported for JK-1. pNKFLAG-5A/del-3 encodes the internally deleted NS5A protein missing aa 235 to 276, a region nonessential for the interaction with NS5B in JK-1. pNKGST-5B/wild encodes the full-length and wild-type NS5B proteins of M1LE, and pNKGST encodes only a GST protein. Under conditions in which the expression levels of FLAG-NS5A proteins (input) were similar and the recovery of the GST-NS5B proteins was almost the same (Fig. 1A, lanes 1 to 5, and C, lanes 2 to 5), coprecipitated NS5A proteins (output) were examined (Fig. 1B, lanes 1 to 5). FLAG-NS5A/wild bound to GST-NS5B (Fig. 1B, lane 2) but not to GST alone (Fig. 1B, lane 1). This result demonstrates that NS5A and NS5B also interact not only in JK-1 but also in M1LE in vivo. Very little FLAG-NS5A/del-1 or /del-2 was recovered (Fig. 1B, lane 3 and 4); however, FLAG-NS5A/del-3 was efficiently pulled down (Fig. 1B, lane 5). When larger amounts of proteins were used for this assay, both FLAG-NS5A/del-1 and /del-2 were weakly detected in the fraction pulled down with GST-NS5B but much significantly weaker than FLAG-5A/wild and /del-3. These results demonstrate that aa 94 to 162 (defined as region 1) and aa 277 to 334 (region 2), but not aa 235 to 276 (region 3), of NS5A seem to be essential for binding NS5B in M1LE as observed in JK-1.

Effect of binding NS5B on HCV RNA replication. To examine the effect of the interaction between NS5A and NS5B on HCV RNA replication in the replicon system, we prepared three kinds of internal deletion mutants, M1LE/5Adel-1, M1LE/5Adel-2, and M1LE/5Adel-3, missing regions 1, 2, and 3 of NS5A, respectively (Fig. 2). M1LE/5Adel-1 and M1LE/5Adel-2 are impaired in their binding to NS5B, but M1LE/5Adel-3 is not. As a negative control, we prepared M1LE/5B-VDD, in which the GDD motif of NS5B was mutated to VDD.

FLAG-tagged wild-type and internally deleted NS5A proteins were efficiently expressed in transiently transfected COS1 cells (Fig. 3). When wild-type M1LE and M1LE/5Adel-3 were transfected by electroporation into our laboratory's Huh-7 cell line, Huh-7-DMB, G418-resistant colonies emerged after selection at a concentration of 1 mg/ml. In the case of M1LE/5Adel-3, the number of G418-resistant colonies was about seven times fewer than in wild-type M1LE. In contrast, no colonies emerged when M1LE/5Adel-1, M1LE/5Adel-2, and M1LE/5B-VDD were transfected into Huh-7-DMB cells, indicating that both of the NS5B-interacting regions of NS5A are critical for HCV RNA replication (Fig. 4). The fact that M1LE/5Adel-3 was replication competent but less efficient than the wild type in Huh-7-DMB cells may reflect some roles of region 3 in HCV RNA replication or a conformational change introduced by the internal deletion (see Discussion).

To minimize the effect of the internal deletion and further delineate the sequence(s) critical for HCV RNA replication, we used the alanine-scanning method (7). All residues of the two regions were scanned by introducing alanine substitution




FIG. 1. Interaction between NS5A and NS5B of the isolate HCV M1LE and the regions essential for this interaction. COS1 cells were transiently cotransfected with mammalian expression vectors expressing FLAG-NS5A proteins (lanes: 1 and 2, wild type; 3, internal deletion mutant 1; 4, internal deletion mutant 2; 5, internal deletion mutant 3) and GST protein alone (lane 1) or GST-NS5B proteins (lanes 2, 3, 4, and 5). (A) Input of FLAG-NS5A proteins. Total lysates were fractionated by SDS-10% PAGE and subjected to Western blot analysis with anti-FLAG monoclonal antibody. (B) Output of FLAG-NS5A proteins. Coprecipitants by glutathione resin were washed with PBS (–) containing 1.0% Triton X-100, fractionated by SDS-10% PAGE, and subjected to Western blot analysis with anti-FLAG monoclonal antibody. (C) Recovery of GST or GST-NS5B proteins. The nitrocellulose membrane used for Western blot analysis of coprecipitants with anti-FLAG antibody was reprobed with anti-GST antibody. Molecular masses (in kilodaltons) are indicated to the right of each panel.

cm in addition to one cm mutant within region 3 as a control (Fig. 2). All of these cm mutants in FLAG-tagged forms, 10 in region 1, 9 in region 2, and 1 in region 3, were similarly expressed in transiently transfected COS1 cells (Fig. 3). When these 20 mutants were transfected by electroporation into Huh-7-DMB cells, only M1LE/cm 252, 277, 283, 290, 297, and 304 were found to be replication competent, although less so than wild-type M1LE. All other mutants were replication incompetent (Fig. 5). The regions of cm 252, 277, 283, 290, 297, and 304 are predicted to form a helical structure by DNASIS-Mac, version 3.2 (Hitachi Software Engineering Co.). The competence of replication may be due to this original structure (see Discussion). To rule out this possibility, we constructed another cm mutant, M1LE/cm 110. In this mutant, the region from aa 110 to 117 of NS5A is predicted to form a helical structure, were all changed to alanines, and after the transfection into Huh-7-DMB cells, no colonies emerged (data not shown). These results support the notion that the inability of the internal deletion mutants, M1LE/5Adel-1 and M1LE/5Adel-2, to replicate is due not to conformational change induced by the deletions but to the absence of interaction between NS5A and NS5B. It is also unlikely that these results are due to an increased cytotoxicity associated with the mutant NS5As, because we observed no decrease in transfection efficiency or ability to establish colonies by using the plasmid

encoding a drug resistance marker along with the wild or the mutant NS5A protein (data not shown).

Improvement in the HCV replicon system. The results clearly showed that two discontinuous regions of NS5A are essential for HCV RNA replication by using the HCV replicon system with Huh-7-DMB cells; however, the number of G418-resistant colonies per microgram of transfected RNA was much smaller than previously reported (24, 29, 35, 37). It remains unclear whether some mutants were replication competent but too inefficient to be detected in the system we applied. Therefore, we tried to improve the assay system in two ways, by the introduction of point mutations to NS5A and by the selection of Huh-7 cells cured of HCV RNA replication by IFN treatment.

We constructed three mutants, M1LE/S225P, M1LE/delS229, and M1LE/S232I, harboring the point mutation S225P (35), a deletion of S229 (delS229) (24), and S232I (4), respectively, all defined as adaptive mutations in other HCV replicon systems (Fig. 2). Next, the 50-1 cells, an HCV subgenome-replicating subclone, were cured of HCV RNA by treatment with IFN for 2 weeks (as described in Materials and Methods), and then the absence of HCV RNA was determined from the results of Northern hybridization, reverse transcription-PCR, and sensitivity to G418 (data not shown). The 50-1 cells cured of HCV RNA by treatment with IFN, designated




FIG. 2. (A) Schematic presentation of HCV mutant replicons used in this study. Wild-type M1LE contains the HCV M1LE wild-type sequence. M1LE/5Adel-1, M1LE/5Adel-2, and M1LE/5Adel-3 encode internal deletions of NS5A missing aa 94 to 162 (region 1), aa 277 to 334 (region 2), and aa 235 to 276 (region 3), respectively. All amino acids within region 1 were replaced with alanines, and 11 cm mutants were prepared. All amino acids within region 2 were replaced, and 9 cm mutants were prepared. The aa 252 to 258 within region 3 were replaced, and M1LE/cm 252 was prepared. The point mutation S225P, a deletion of S229, and S232I were introduced into wild-type M1LE, and then M1LE/S225P, M1LE/delS229, and M1LE/S232I were prepared. M1LE/5B-VDD encodes NS5B in which the GDD motif was mutated to VDD. (B) Summary of the NS5A mutations. This figure shows the entire amino acid sequence of NS5A of M1LE and the positions of mutations. Numbering starts from the beginning of NS5A. Internal deletions 1, 2, and 3, and point mutations S225P, delS229, and S232I, and cm 252 were introduced as described for panel A. All amino acids within region 1 were replaced with alanines, and then M1LE/cm 94, 100, 105, 110, 113, 120, 127, 134, 141, 148, and 155 were prepared. All amino acids within region 2 were replaced with alanines, and M1LE/cm 277, 283, 290, 297, 304, 311, 316, 321, and 328 were prepared. The positions of the substituted amino acids in each cm mutant are shown in panel B.




FIG. 3. Efficient translation of mutant FLAG-NS5As. COS1 cells were transiently transfected with mammalian expression vectors expressing FLAG-NS5A proteins that were prepared as described in Materials and Methods. Lanes: 4, 16, and 26, wild type; 1, internal deletion 1; 2, internal deletion 2; 3, internal deletion 3; 5, cm 252; 6, cm 94; 7, cm 100; 8, cm 105; 9, cm 113; 10, cm 120; 11, cm 127; 12, cm 134; 13, cm 141; 14, cm 148; 15, cm 155; 17, cm 277; 18, cm 283; 19, cm 290; 20, cm 297; 21, cm 304; 22, cm 311; 23, cm 315; 24, cm 321; 25, cm 328. Total lysates were fractionated by SDS-10% PAGE and subjected to Western blot analysis with anti-FLAG monoclonal antibody. Molecular masses (in kilodaltons) are indicated to the left of the panels.

Huh-7-KV-C, were evaluated. Wild-type and these mutant HCV replicons were transfected into Huh-7-DMB and Huh-7-KV-C cells. In Huh-7-DMB cells, M1LE/S225P and M1LE/S232I actually increased the efficiency of transduction to some extent, but M1LE/delS229 did not (Fig. 6A). M1LE/S232I was the most effective. In Huh-7-KV-C cells, interestingly, no colonies emerged after selection with G418 when wild-type M1LE was transfected, whereas in the case of M1LE/S225P, delS229, and S232I, 2,500, 3,000, and 25,000 colonies/ μ g of RNA emerged, respectively (Fig. 6B). These results indicate that two point mutations, S225P and S232I, can be categorized as the adaptive mutations in the isolate M1LE and that the cells cured of HCV RNA by treatment with IFN, Huh-7-KV-C, show higher permissiveness for M1LE/S225P, delS229, and S232I than Huh-7-DMB cells. In this way, we established highly improved replicon systems.

Delineation of important sequences of NS5A for HCV RNA replication. To examine the effect of internal deletions and alanine substitutions on HCV RNA replication with this improved replicon system, double mutants with S232I plus internal deletion mutations or alanine-substituted cm's in M1LE were constructed and transfected into Huh-7-DMB and Huh-7-KV-C cells by electroporation. After G418 selection, in Huh-7-DMB cells, some 400 to 1,000 colonies/ μ g of RNA emerged with the double mutants of M1LE/S232I plus cm 252, 277, 283, 297, and 304, but only about 100 colonies/ μ g of RNA emerged with the double mutant M1LE/S232I plus cm 290. No colonies

emerged with the double mutant M1LE/S232I plus other cm's, del-1, del-2, and del-3 (Fig. 7; data not shown for Huh-7-KV-C cells). The double mutant M1LE/S232I plus cm 110 was also replication incompetent in Huh-7-DMB cells. To further examine the replication competence of these mutants, double mutants of M1LE/S225P plus an internal deletion mutation or cm were constructed and then transfected into Huh-7-DMB cells. The results were almost the same as those with the double mutants with S232I, although the efficiencies of these mutants were around one-half of those with S232I (Fig. 8). The similar results in the double mutants with S225P were obtained with Huh-7 KV-C cells (data not shown). The replication-defective property of the mutants with deletions of regions 1 and 2, the cm mutants within region 1, and the cm mutants in the C-terminal part of region 2 was also observed with both Huh-7-DMB and Huh-7-KV-C cells, indicating that the replication incompetence of these mutants is not due to the low efficiency of the original assay system. The cm mutants at the N-terminal part of region 2, cm 277, 283, 290, 297, and 304, were replication competent in the absence of the adaptive mutation in Huh-7-DMB cells and also in the presence of the adaptive mutation in Huh-7-DMB and Huh-7-KV-C cells to some extent. Taken together, these results suggest that the interaction with NS5B through regions 1 and 2, probably through its C-terminal part, is also essential for HCV RNA replication. We examined the interaction between cm mutants of FLAG-NS5A and GST-NS5B, but the difference among wild-type and cm mutants was weak in the pull-down assay. Differential binding would be possible if the two partner proteins were lower in concentration or together with other NS proteins, as those occur in vivo in HCV-RNA-replicating cells.

Interestingly, two quantitative differences were observed with M1LE/5Adel-3, cm 252, and 290 with and without the adaptive mutations. M1LE/5Adel-3 was weakly replication competent in Huh-7-DMB cells but incompetent in Huh-7-DMB and Huh-7-KV-C cells when the adaptive mutations were introduced. In contrast, M1LE/cm 252 was weakly replication competent in Huh-7-DMB cells but as high as that of the other replication-competent cm mutants in the presence of the adaptive mutations in Huh-7-DMB and Huh-7-KV-C cells (data not shown for Huh-7-KV-C cells; see Discussion).

DISCUSSION

HCV NS5A is a viral regulatory protein that modulates viral RNA replication and host processes by interacting directly and indirectly with a variety of host regulatory factors (10, 19, 42, 56, 58, 67). The important role of NS5A in HCV RNA replication has been clearly demonstrated by high or clustered incidence of adaptive mutations in NS5A detected in HCV RNA replicon systems, although the molecular mechanism involved remains unknown (4, 24, 35, 37, 40). Shirota et al. previously reported direct interaction between NS5A and NS5B through two binding regions of NS5A expressed in mammalian cells and in vitro with a purified recombinant and that NS5A could modulate the activity of NS5B RdRp in vitro through this direct interaction (54). Here we demonstrated the critical role of regions essential for the NS5A-NS5B interaction in HCV RNA replication with an HCV subgenomic replicon by introducing several internal deletion mutations into




FIG. 4. Effect of internal deletion mutation on HCV RNA replication. Huh-7-DMB cells were transfected with 1 μ g of in vitro-transcribed wild-type M1LE, 5Adel-1, 5Adel-2, 5Adel-3, and 5B-VDD RNA by electroporation, and G418-resistant cells were selected with a G418 concentration of 1 mg/ml. G418-resistant cell colonies were stained 4 weeks after transfection. (A) This figure shows the mean number of G418-resistant cell colonies isolated per 10-cm-diameter cell culture dish per 1 μ g of RNA. Error bars indicate the standard deviations of the results from at least three independent experiments. (B) G418-resistant colonies were visualized by staining cells as described in Materials and Methods.

NS5A. Our results provide clear evidence that NS5A is indispensable for HCV RNA replication probably through its interaction with NS5B, since the mutants missing the NS5B-binding regions, regions 1 and 2, could not produce G418-resistant colonies, but the mutants missing region 3, which is not essential for this interaction, could. The critical role of NS5A in HCV RNA replication in this report is consistent with the previous one that an amino-terminal amphipathic α -helix of NS5A is essential for HCV RNA replication in the replicon system, localizing NS5A to a membrane (13). The use of cm mutants strongly suggests that all of region 1 and the C-terminal part of region 2 are critical for HCV RNA replication. However, it is difficult to exclude the possibility that the substitution of eight amino acids in a row may induce structural change. Then the critical regions defined by cm mutants may simply reflect the structural integrity necessary for the function, although a structural evaluation is difficult at present, since no crystal model of NS5A is available. Further mutational analysis is necessary to test these possibilities.

HCV RNA replication would take place in a distinctly altered membrane structure of the endoplasmic reticulum, a membranous web (12), as recently reported by Gosert et al. (20). All NS proteins might be recruited to the membrane

structure via their own membrane association domains or by the help of NS4A in the case of NS3 (8, 13, 20, 28, 30, 51, 52, 63). Recently, Dimitrova et al. (11) reported that all six NS proteins interact with each other through their multiple interacting surfaces. NS5B is HCV RdRp and has been reported to interact with NS proteins and some host proteins. Such interaction(s) may modulate the activity of NS5B RdRp in various ways. The critical role of the homomeric interaction of NS5B in RdRp activity was demonstrated by us and another group (48, 60). Piccininni et al. (47) reported that NS5B interacts with NS3 and NS4B as positive and negative regulators in the replication complex. Previously, it was reported that the direct binding of NS5A and NS5B in the isolate JK-1 weakly stimulated the activity of NS5B RdRp in vitro at first (at a molar ratio to NS5B of less than 0.1) and then inhibited the activity in a dose-dependent manner (54). In the present study, we showed that the two regions of NS5A are important for binding NS5B and are essential for HCV RNA replication in the isolate M1LE by HCV replicon assays. The weak stimulation by NS5A of RdRp activity through the binding of NS5B observed in vitro may reflect the essential role of NS5A in HCV RNA replication, or the interaction between NS5A and NS5B is important for the dynamic assembly of NS proteins in the HCV




FIG. 5. Effect of clustered alanine-substitution mutations on HCV RNA replication. Huh-7-DMB cells were transfected with 1 µg of in vitro-transcribed wild-type M1LE, 5Adel-1, 5Adel-2, and cm 94, 100, 105, 113, 120, 127, 134, 141, 148, 155, 252, 277, 283, 290, 297, 304, 311, 316, 321, and 328 RNA, and G418-resistant cells were selected with a G418 concentration of 0.5 mg/ml. G418-resistant cell colonies were stained 4 weeks after transfection. This figure shows the mean number of G418-resistant cell colonies isolated per 15-cm-diameter cell culture dish per 1 µg of RNA. Error bars indicate the standard deviations of the results from at least three independent experiments.




FIG. 6. Effects of three kinds of point mutation, S225P, deletion S229, and S232I, on HCV RNA replication in the Huh-7-DMB and KV-C sublines. 50-1 cells were cured of self-replicating subgenomic RNAs by IFN-α treatment, and then cured 50-1 cells, Huh-7-KV-C, were prepared. Huh-7-DMB and KV-C cells were transfected with 10 ng to 1 µg of in vitro-transcribed wild-type M1LE, M1LE/S225P, M1LE/delS229, and M1LE/S232I RNA. G418-resistant cells were selected with a G418 concentration of 1 mg/ml. G418-resistant cell colonies were stained 4 weeks after transfection. (A) This figure shows the mean number of G418-resistant cell colonies isolated per 10-cm-diameter cell culture dish per 1 µg of RNA when each in vitro-transcribed mutant RNA was transfected into Huh-7-DMB cells. Error bars indicate the standard deviations of the results from at least three independent experiments. (B) This figure shows the mean number of G418-resistant cell colonies isolated per 10-cm-diameter cell culture dish per 1 µg of RNA when each in vitro-transcribed mutant RNA was transfected into Huh-7-KV-C cells. Error bars indicate the standard deviations of the results from at least three independent experiments.




FIG. 7. Effect of M1LE/S232I plus internal deletion mutations or alanine-substitution mutations. Huh-7-DMB cells were transfected with 1 μ g of in vitro-transcribed M1LE/S232I plus 5Adel-1, 5Adel-2, 5Adel-3, and cm 94, 100, 105, 113, 120, 127, 134, 141, 148, 155, 252, 277, 283, 290, 297, 304, 311, 316, 321, and 328 RNA. G418-resistant cells were selected with a G418 concentration of 1 mg/ml and stained 4 weeks after transfection. This figure shows the mean number of G418-resistant cell colonies isolated per 10-cm-diameter cell culture dish per 1 μ g of RNA. Error bars indicate the standard deviations of the results from at least three independent experiments. The number of G418-resistant cell colonies in M1LE/S232I is derived from the results shown in Fig. 6A.

replication complex. Alternatively, it cannot be excluded that the internal deletion and cm mutants may be defective in other unknown function(s) essential for HCV RNA replication.

Adaptive mutations that increase the efficiency of HCV RNA replication have been accumulated in different HCV replicon systems (4, 24, 35, 37, 40). We introduced several point mutations into NS5A of M1LE and found that two, S232I and S225P, positively affected colony formation as adaptive mutations in two different Huh-7 sublines. Interestingly, a deletion of S229 had an effect distinct from those of the other two

mutations, since it was only effective in the cured cells, the KV-C subline. Also, it is noteworthy that wild-type M1LE could not replicate in the Huh-7-KV-C subline. The effect of the deletion of S229 and the replication incompetence of wild-type M1LE may be of interest for the elucidation of the phenotypic or genetic change(s) in the cured cells. In combination with the adaptive mutation, the deletion and cm mutants exhibited distinct phenotypes in HCV RNA replication.

First, the mutant missing region 3 was replication incompetent in the presence of S232I or S225P, which was in contrast




FIG. 8. Effect of M1LE/S225P plus internal deletion mutations or alanine-substitution mutations. Huh-7-DMB cells were transfected with 1 μ g of in vitro-transcribed M1LE/S225P plus 5Adel-2, 5Adel-3, and cm 252, 277, 283, 290, 297, 304, 311, 316, 321, and 328 RNA. G418-resistant cells were selected with a G418 concentration of 1 mg/ml and stained 4 weeks after transfection. This figure shows the mean number of G418-resistant cell colonies isolated per 10-cm-diameter cell culture dish per 1 μ g of RNA. Error bars indicate the standard deviations of the results from at least three independent experiments. The number of G418-resistant cell colonies in M1LE/S225P is derived from the results shown in Fig. 6A.

to the dramatic increase in HCV RNA replication efficiency obtained by introducing the adaptive mutation alone. A similar but more moderate phenotype was observed with cm 252 when the adaptive mutation was present. Our result is different from the reported enhancement of G418-resistant colony formation caused by a large 47-aa deletion in HCV-Con1 (4) and by a natural 4-aa insertion in HCV-N (29). Region 3 contains the IFN sensitivity-determining region, within which mutations have been reported to be associated with sensitivity to IFN therapy in patients with chronic HCV infection in Japan (14, 15). Region 3 is almost the same as a protein kinase R (PKR)-binding domain (18). It has been reported that NS5A can disrupt the dimerization of PKR through binding, resulting in the repression of PKR function, and efficient HCV RNA replication may involve a block in PKR-dependent signaling (18, 46). In this context, the adaptive mutations can greatly augment HCV RNA replication and thus may induce PKR, which could be inactivated via interaction with the PKR-binding region of NS5A, region 3, but not with those mutants defective in PKR-binding, such as del-3 and cm 252.

Second, cm 290 was less replication competent than the other cm mutants in region 2 when the adaptive mutation, S232I or S225P, was present. The highly charged sequence mutated in cm 290 may be critical for HCV RNA replication in the presence of the adaptive mutation. To address this point, four different combinations of three to four alanine substitutions in 7 aa residues were introduced into the M1LE/S232I construct. All of these mutants were more efficient in HCV RNA replication than M1LE/S232I plus cm 290 but still less so than the other replication-competent mutants in region 2 (data not shown), suggesting that all or most amino acids in the sequence contribute to its critical role in HCV RNA replication in the presence of the adaptive mutations. This result may suggest a functional linkage of the sequence mutated in cm 290 to the adaptive mutations.

In HCV subgenomic replicons, some groups have found that cured cell clones showed a high permissiveness for HCV RNA replication (5, 45) while another has not (40). Between the recipient sublines we used, the cured cells (Huh-7-KV-C) shared the nonpermissive property for wild-type M1LE; however, the DMB subline was permissive for wild-type M1LE and/or adaptive mutations with lower efficiency than the KV-C subline. These results suggest that several different genetic or phenotypic alterations in recipient cells emerge under IFN treatment or multiple ways for cells to be permissive to HCV RNA replication.

In summary, we established a highly efficient HCV replicon system derived from the isolate M1LE and demonstrated that the two regions critical for the interaction between NS5A and NS5B are also indispensable for HCV RNA replication in an HCV replicon system. Our results strongly suggest that NS5A is involved in the HCV replication complex and acts as a positive modulator of HCV RNA replication through its interaction with NS5B. The molecular mechanism of this positive effect by NS5A remains to be elucidated and may lead to the design of new drugs that inhibit HCV RNA replication.

ACKNOWLEDGMENTS

We are grateful to N. Hayashi and T. Nomura for encouraging discussion. We also thank M. Yasukawa and K. Kuwabara for technical assistance.

This work was supported in part by the Program for Promotion of Fundamental Studies in Health Sciences of the Organization for Pharmaceutical Safety and Research, Grants-in-aid for scientific research (B) and development, and Grants-in-aid for scientific research on priority areas (C) in oncogenesis from the Ministry of Education, Sports, Culture, and Technology.

REFERENCES

- Alter, H. J., R. H. Purcell, J. W. Shih, J. C. Melpolder, M. Houghton, Q. L. Choo, and G. Kuo. 1989. Detection of antibody to hepatitis C virus in prospectively followed transfusion recipients with acute and chronic non-A, non-B hepatitis. *N. Engl. J. Med.* **321**:1494–1500.
- Bartenschlager, R., L. Ahlborn-Laake, J. Mous, and H. Jacobsen. 1993. Nonstructural protein 3 of the hepatitis C virus encodes a serine-type proteinase required for cleavage at the NS3/4 and NS4/5 junctions. *J. Virol.* **67**:3835–3844.
- Behrens, S. E., L. Tomei, and R. De Francesco. 1996. Identification and properties of the RNA-dependent RNA polymerase of hepatitis C virus. *EMBO J.* **15**:12–22.
- Blight, K. J., A. A. Kolykhalov, and C. M. Rice. 2000. Efficient initiation of HCV RNA replication in cell culture. *Science* **290**:1972–1974.
- Blight, K. J., J. A. McKeating, and C. M. Rice. 2002. Highly permissive cell lines for subgenomic and genomic hepatitis C virus RNA replication. *J. Virol.* **76**:13001–13014.
- Blight, K. J., J. A. McKeating, J. Marcotrigiano, and C. M. Rice. 2003. Efficient replication of hepatitis C virus genotype 1a RNAs in cell culture. *J. Virol.* **77**:3181–3190.
- Bordo, D., and P. Argos. 1991. Suggestion for safe residue substitutions in site-directed mutagenesis. *J. Mol. Biol.* **217**:721–729.
- Brass, V., E. Bieck, R. Montserret, B. Wölk, J. A. Hellings, H. E. Blum, F. Penin, and D. Moradpour. 2002. An amino-terminal amphipathic α -helix mediates membrane association of the hepatitis C virus nonstructural protein 5A. *J. Biol. Chem.* **277**:8130–8139.
- Choo, Q. L., G. Kuo, A. J. Weiner, L. R. Overby, D. W. Bradley, and M. Houghton. 1989. Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* **244**:359–362.
- Chung, K. M., J. Lee, J. E. Kim, O. K. Song, S. Cho, J. Lim, M. Seedorf, B. Hahn, and S. K. Jang. 2000. Nonstructural protein 5A of hepatitis C virus inhibits the function of karyopherin β 3. *J. Virol.* **74**:5233–5241.
- Dimirova, M., I. Imbert, M. P. Kieney, and C. Schuster. 2003. Protein-protein interactions between hepatitis C virus nonstructural proteins. *J. Virol.* **77**:5401–5414.
- Egger, D., B. Wölk, R. Gosert, L. Bianchi, H. E. Blum, D. Moradpour, and K. Bienz. 2002. Expression of hepatitis C virus proteins induces distinct membrane alterations including a candidate viral replication complex. *J. Virol.* **76**:5974–5984.
- Elazar, M., K. H. Cheong, P. Liu, H. B. Greenberg, C. M. Rice, and J. S. Glenn. 2003. Amphipathic helix-dependent localization of NS5A mediates hepatitis C virus RNA replication. *J. Virol.* **77**:6055–6061.
- Enomoto, N., I. Sakuma, Y. Asahina, M. Kurosaki, T. Murakami, C. Yamamoto, N. Izumi, F. Marumo, and C. Sato. 1995. Comparison of full-length sequence of interferon-sensitive and resistant hepatitis C virus 1b. Sensitivity to interferon is conferred by amino acid substitutions in the NS5A region. *J. Clin. Invest.* **96**:224–230.
- Enomoto, N., I. Sakuma, Y. Asahina, M. Kurosaki, T. Murakami, C. Yamamoto, Y. Ogura, N. Izumi, F. Marumo, and C. Sato. 1996. Mutations in the nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N. Engl. J. Med.* **334**:77–82.
- Ferrari, E., J. Wright-Minogue, J. W. S. Fang, B. M. Baroudy, J. Y. N. Lau, and Z. Hong. 1999. Characterization of soluble hepatitis C virus RNA-dependent RNA polymerase expressed in *Escherichia coli*. *J. Virol.* **73**:1649–1654.
- Friebe, P., and R. Bartenschlager. 2002. Genetic analysis of sequence in the 3' nontranslated region of hepatitis C virus that are important for RNA replication. *J. Virol.* **76**:5326–5338.
- Gale, M., Jr., C. M. Blakely, B. Kwieciszewski, S. L. Tan, M. Dossett, N. M. Tang, M. J. Korth, S. J. Polyak, D. R. Gretch, and M. G. Katze. 1998. Control of PKR protein kinase by hepatitis C virus nonstructural 5A protein: molecular mechanisms of kinase regulation. *Mol. Cell. Biol.* **18**:5208–5218.
- Ghosh, A. K., M. Majumder, R. Steele, P. Yaciuk, J. Chivria, R. Ray, and R. B. Ray. 2000. Hepatitis C virus NS5A protein modulates transcription through a novel cellular transcription factor SRCAP. *J. Biol. Chem.* **275**:7184–7188.
- Gosert, R., D. Egger, V. Lohmann, R. Bartenschlager, H. E. Blum, K. Bienz, and D. Moradpour. 2003. Identification of the hepatitis C virus RNA replication complex in Huh-7 cells harboring subgenomic replicons. *J. Virol.* **77**:5487–5492.
- Grakoui, A., D. W. McCourt, C. Wychowski, S. M. Feinstone, and C. M. Rice. 1993. Characterization of the hepatitis C virus-encoded serine protease: determination of proteinase-dependent polyprotein cleavage sites. *J. Virol.* **67**:2832–2843.
- Grakoui, A., C. Wychowski, C. Lin, S. M. Feinstone, and C. M. Rice. 1993.

- Expression and identification of hepatitis C virus polyprotein cleavage products. *J. Virol.* **67**:1385–1395.
23. Gu, B., A. T. Gates, O. Isken, S. E. Behrens, and R. T. Sarisky. 2003. Replication studies using genotype 1a subgenomic hepatitis C virus replicons. *J. Virol.* **77**:5352–5359.
 24. Guo, J. T., V. V. Bichko, and C. Seeger. 2001. Effect of alpha interferon on the hepatitis C virus replicon. *J. Virol.* **75**:8516–8523.
 25. Hijikata, M., H. Mizushima, Y. Tanji, Y. Komoda, Y. Hirowatari, T. Akagi, N. Kato, K. Kimura, and K. Shimotohno. 1993. Proteolytic processing and membrane association of putative nonstructural proteins of hepatitis C virus. *Proc. Natl. Acad. Sci. USA* **90**:10773–10777.
 26. Hijikata, M., N. Kato, Y. Ootsuyama, M. Nakagawa, and K. Shimotohno. 1991. Gene mapping of the putative structural region of the hepatitis C virus genome by *in vitro* processing analysis. *Proc. Natl. Acad. Sci. USA* **88**:5547–5551.
 27. Hijikata, M., N. Kato, Y. Ootsuyama, M. Nakagawa, S. Ohkoshi, and K. Shimotohno. 1991. Hypervariable regions in the putative glycoprotein of hepatitis C virus. *Biochem. Biophys. Res. Commun.* **175**:220–228.
 28. Hügler, T., F. Fehrmann, E. Bieck, M. Kohara, H. G. Kräusslich, C. M. Rice, H. E. Blum, and D. Moradpour. 2001. The hepatitis C virus nonstructural protein 4B is an integral endoplasmic reticulum membrane protein. *Virology* **284**:70–81.
 29. Ikeda, M., M. Yi, K. Li, and S. M. Lemon. 2002. Selectable subgenomic and genome-length dicistronic RNAs derived from an infectious molecular clone of the HCV-N strain of hepatitis C virus replicate efficiently in cultured Huh7 cells. *J. Virol.* **76**:2997–3006.
 30. Ivashkina, N., B. Wölk, V. Lohmann, R. Bartenschlager, H. E. Blum, F. Penin, and D. Moradpour. 2002. The hepatitis C virus RNA-dependent RNA polymerase membrane insertion sequence is a transmembrane segment. *J. Virol.* **76**:13088–13093.
 31. Kato, N., T. Nakazawa, T. Mizutani, and K. Shimotohno. 1995. Susceptibility of human T-lymphotropic virus type I-infected cell line MT-2 to hepatitis C virus infection. *Biochem. Biophys. Res. Commun.* **206**:863–869.
 32. Kato, N., Y. Ootsuyama, H. Sekiya, S. Ohkoshi, T. Nakazawa, M. Hijikata, and K. Shimotohno. 1994. Genetic drift in hypervariable region 1 of the viral genome in persistent hepatitis C virus infection. *J. Virol.* **68**:4776–4784.
 33. Kishine, H., K. Sugiyama, M. Hijikata, N. Kato, H. Takahashi, T. Noshi, Y. Nio, M. Hosaka, Y. Miyazaki, and K. Shimotohno. 2002. Subgenomic replicon derived from a cell line infected with the hepatitis C virus. *Biochem. Biophys. Res. Commun.* **293**:993–999.
 34. Kolykhalov, A. A., K. Mihalik, S. M. Feinstone, and C. M. Rice. 2000. Hepatitis C virus-encoded enzymatic activities and conserved RNA elements in the 3' nontranslated region are essential for virus replication *in vivo*. *J. Virol.* **74**:2046–2051.
 35. Krieger, N., V. Lohmann, and R. Bartenschlager. 2001. Enhancement of hepatitis C virus RNA replication by cell culture-adaptive mutations. *J. Virol.* **75**:4614–4624.
 36. Lin, Y., H. Tang, T. Nomura, D. Dorjsuren, N. Hayashi, W. Wei, T. Ohta, R. Roeder, and S. Murakami. 1998. The hepatitis B virus X protein is a co-activator of activated transcription that modulates the transcription machinery and distal binding activators. *J. Biol. Chem.* **273**:27097–27103.
 37. Lohmann, V., F. Körner, A. Dobierzewska, and R. Bartenschlager. 2001. Mutations in hepatitis C virus RNAs conferring cell culture adaptation. *J. Virol.* **75**:1437–1449.
 38. Lohmann, V., F. Körner, J. O. Koch, U. Herian, L. Theilmann, and R. Bartenschlager. 1999. Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science* **285**:110–113.
 39. Lohmann, V., F. Körner, U. Herian, and R. Bartenschlager. 1997. Biochemical properties of hepatitis C virus NS5B RNA-dependent RNA polymerase and identification of amino acid sequence motifs essential for enzymatic activity. *J. Virol.* **71**:8416–8428.
 40. Lohmann, V., S. Hoffmann, U. Herian, F. Penin, and R. Bartenschlager. 2003. Viral and cellular determinants of hepatitis C virus RNA replication in cell culture. *J. Virol.* **77**:3007–3019.
 41. Major, M. E., and S. M. Feinstone. 1997. The molecular virology of hepatitis C. *Hepatology* **25**:1527–1538.
 42. Majumder, M., A. K. Ghosh, R. Steele, R. Ray, and R. B. Ray. 2001. Hepatitis C virus NS5A physically associates with p53 and regulates p21/waf1 gene expression in a p53-dependent manner. *J. Virol.* **75**:1401–1407.
 43. Mizutani, T., N. Kato, S. Saito, M. Ikeda, K. Sugiyama, and K. Shimotohno. 1996. Characterization of hepatitis C virus replication in cloned cells obtained from a human T-cell leukemia virus type I-infected cell line, MT-2. *J. Virol.* **70**:7219–7223.
 44. Murphy, F. A., C. M. Fauquet, D. H. L. Bishop, S. A. Ghabrial, A. W. Jarvis, G. P. Martelli, M. A. Mayo, and M. D. Summers. 1995. Classification and nomenclature of viruses: sixth report of the international committee on taxonomy of viruses, p. 424–426. Springer-Verlag, Vienna, Austria.
 45. Murray, E. M., J. A. Grobler, E. J. Markel, M. F. Pagnoni, G. Paonessa, A. J. Simon, and O. A. Flores. 2003. Persistent replication of hepatitis C virus replicons expressing the β -lactamase reporter in subpopulations of highly permissive Huh7 cells. *J. Virol.* **77**:2928–2935.
 46. Pflugheber, J., B. Fredericksen, R. Sumpter, Jr., C. Wang, F. Ware, D. L. Sodora, and M. Gale, Jr. 2002. Regulation of PKR and IRF-1 during hepatitis C virus RNA replication. *Proc. Natl. Acad. Sci. USA* **99**:4650–4655.
 47. Piccinini, S., A. Varaklioti, M. Nardelli, B. Dave, K. D. Raney, and J. E. G. McCarthy. 2002. Modulation of the hepatitis C virus RNA-dependent RNA polymerase activity by the non-structural (NS) 3 helicase and the NS4B membrane protein. *J. Biol. Chem.* **277**:45670–45679.
 48. Qin, W., H. Luo, T. Nomura, N. Hayashi, T. Yamashita, and S. Murakami. 2002. Oligomeric interaction of hepatitis C virus NS5B is critical for catalytic activity of RNA-dependent RNA polymerase. *J. Biol. Chem.* **277**:2132–2137.
 49. Qin, W., T. Yamashita, Y. Shiota, Y. Lin, W. Wei, and S. Murakami. 2001. Mutational analysis of the structure and functions of hepatitis C virus RNA-dependent RNA polymerase. *Hepatology* **33**:728–737.
 50. Saito, I., T. Miyamura, A. Ohbayashi, H. Harada, T. Katayama, S. Kikuchi, Y. Watanabe, S. Koi, M. Onji, Y. Ohta, C. Choo, M. Houghton, and G. Kuo. 1990. Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc. Natl. Acad. Sci. USA* **87**:6547–6549.
 51. Santolini, E., L. Pacini, C. Fipaldini, G. Migliaccio, and N. Monica. 1995. The NS2 protein of hepatitis C virus is a transmembrane polypeptide. *J. Virol.* **69**:7461–7471.
 52. Schmidt-Mende, J., E. Bieck, T. Hügler, F. Penin, C. M. Rice, H. E. Blum, and D. Moradpour. 2001. Determinants for membrane association of the hepatitis C virus RNA-dependent RNA polymerase. *J. Biol. Chem.* **276**:44052–44063.
 53. Seeff, L. B. 1997. Natural history of hepatitis C. *Hepatology* **26**:21S–28S.
 54. Shiota, Y., H. Luo, W. Qin, S. Kaneko, T. Yamashita, K. Kobayashi, and S. Murakami. 2002. Hepatitis C virus (HCV) NS5A binds RNA-dependent RNA polymerase (RdRp) NS5B and modulates RNA-dependent RNA polymerase activity. *J. Biol. Chem.* **277**:11149–11155.
 55. Sugiyama, K., N. Kato, T. Mizutani, M. Ikeda, T. Tanaka, and K. Shimotohno. 1997. Genetic analysis of the hepatitis C virus (HCV) genome from HCV-infected human T cells. *J. Gen. Virol.* **78**:329–336.
 56. Tan, S. L., H. Nakao, Y. He, S. Vijaysri, P. Neddermann, B. L. Jacobs, B. J. Mayer, and M. G. Katze. 1999. NS5A, a nonstructural protein of hepatitis C virus, binds growth factor receptor-bound protein 2 adaptor protein in a Src homology 3 domain/ligand-dependent manner and perturbs mitogenic signaling. *Proc. Natl. Acad. Sci. USA* **96**:5533–5538.
 57. Tsukiyama-Kohara, K., N. Iizuka, M. Kohara, and A. Nomoto. 1992. Internal ribosome entry site within hepatitis C virus RNA. *J. Virol.* **66**:1476–1483.
 58. Tu, H., L. Gao, S. T. Shi, D. R. Taylor, T. Yang, A. K. Mircheff, Y. Wen, A. E. Gorbalenya, S. B. Hwang, and M. M. C. Lai. 1999. Hepatitis C virus RNA polymerase and NS5A complex with a SNARE-like protein. *Virology* **263**:30–41.
 59. Wang, C., P. Sarnow, and A. Siddiqui. 1993. Translation of human hepatitis C virus RNA in cultured cells is mediated by internal ribosome-binding mechanism. *J. Virol.* **67**:3338–3344.
 60. Wang, Q. M., M. A. Hockman, K. Staschke, R. B. Johnson, K. A. Case, J. Lu, S. Parsons, F. Zhang, R. Rathnachalam, K. Kirkegaard, and J. M. Colacino. 2002. Oligomerization and cooperative RNA synthesis activity of hepatitis C virus RNA-dependent RNA polymerase. *J. Virol.* **76**:3865–3872.
 61. Weiner, A. J., H. M. Geysen, C. Christopherson, J. E. Hall, T. J. Mason, G. Saracco, F. Bonino, K. Crawford, C. D. Marion, K. A. Crawford, M. Brunetto, P. J. Barr, T. Miyamura, J. McHutchinson, and M. Houghton. 1992. Evidence for immune selection of hepatitis C virus (HCV) putative envelope glycoprotein variants: potential role in chronic HCV infections. *Proc. Natl. Acad. Sci. USA* **89**:3468–3472.
 62. Weiner, A. J., M. J. Brauer, J. Rosenblatt, K. H. Richman, J. Tung, K. Crawford, F. Bonino, G. Saracco, Q. L. Choo, M. Houghton, and J. H. Han. 1991. Variable and hypervariable domains are found in the regions of HCV corresponding to the flavivirus envelope and NS1 proteins and the pestivirus envelope glycoproteins. *Virology* **180**:842–848.
 63. Wölk, B., D. Sansonno, H. G. Kräusslich, F. Dammacco, C. M. Rice, H. E. Blum, and D. Moradpour. 2000. Subcellular localization, stability, and trans-cleavage competence of the hepatitis C virus NS3-NS4A complex expressed in tetracycline-regulated cell lines. *J. Virol.* **74**:2293–2304.
 64. Yamashita, T., S. Kaneko, Y. Shiota, W. Qin, T. Nomura, K. Kobayashi, and S. Murakami. 1998. RNA-dependent RNA polymerase activity of the soluble recombinant hepatitis C virus NS5B protein truncated at the C-terminal region. *J. Biol. Chem.* **273**:15479–15486.
 65. Yanagi, M., M. St. Claire, S. U. Emerson, R. H. Purcell, and J. Bukh. 1999. *In vivo* analysis of the 3' untranslated region of the hepatitis C virus after *in vitro* mutagenesis of an infectious cDNA clone. *Proc. Natl. Acad. Sci. USA* **96**:2291–2295.
 66. Yi, M., and S. M. Lemon. 2003. 3' Nontranslated RNA signals required for replication of hepatitis C virus RNA. *J. Virol.* **77**:3557–3568.
 67. Zech, B., A. Kurtenbach, N. Krieger, D. Strand, S. Blencke, M. Morbitzer, K. Salassidis, M. Cotten, J. Wissing, S. Obert, R. Bartenschlager, T. Herget, and H. Daub. 2003. Identification and characterization of amphiphysin II as a novel cellular interaction partner of the hepatitis C virus NS5A protein. *J. Gen. Virol.* **84**:555–560.